

225

by ISST 20205

Submission date: 30-Oct-2023 04:01PM (UTC+0700)

Submission ID: 2211768418

File name: 20205.docx (36.82K)

Word count: 2894

Character count: 16650

Effect of Probiotics Addition on Artificial Feed for catfish Growth

Wiwit Denny Fitriana^{1*}, Bakri², Mukhamad Masrur³, Anna Qomariana⁴, Chandra Sukma Anugrah⁵

¹Faculty of Science and Technology, Universitas Pesantren Tinggi Darul Ulum Jombang, 61481, Jombang, Indonesia

²Faculty of Islamic, Universitas Pesantren Tinggi Darul Ulum Jombang, 61481, Jombang, Indonesia

³Faculty of Science and Technology, Universitas Pesantren Tinggi Darul Ulum Jombang, 61481, Jombang, Indonesia

⁴Faculty of Islamic, Universitas Pesantren Tinggi Darul Ulum Jombang, 61481, Jombang, Indonesia

⁵Faculty of Science and Technology, Universitas Pesantren Tinggi Darul Ulum Jombang, 61481, Jombang, Indonesia

Abstract. The research aimed to investigate the impact of probiotics supplementation in artificial feed on the growth of catfish (*Clarias gariepinus*). Catfish farming is a vital component of aquaculture, and optimizing feed formulations can enhance both the economic and environmental sustainability of the industry. In this study, probiotics were incorporated into the catfish feed to assess their potential benefits on growth performance. A fully randomized design was employed, encompassing four distinct treatment groups denoted as follows: P0, representing the control group without any treatment; P1, involving the incorporation of 2 cc/kg of probiotic into the feed; P2, entailing the inclusion of 4 cc/kg of probiotic in the feed; and P3, comprising the use of 6 cc/kg of probiotic in the feed. Data analysis was conducted using analysis of variance with a confidence level set at 5%. The growth parameters, including weight gain, length increase, and feed conversion ratio (FCR), were monitored over 2 months. The results revealed a significant improvement in the growth performance of catfish in the probiotics-supplemented group compared to the control group. Catfish fed with probiotics exhibited a higher average weight gain and increased length, indicating better growth. Moreover, the FCR was notably improved in the probiotics-treated group, suggesting enhanced feed utilization efficiency. This study highlights the positive effects of probiotics supplementation in artificial feed on the growth of catfish.

Keywords: Probiotic, Fish farming, Catfish, Composition, Aquaculture

1 Introduction

Jombang, located in East Java, is a renowned city known for its thousands of Islamic boarding school students, commonly referred to as "santri." The Millennial Farming Students Movement, which focuses on freshwater fish farming, is one of the programs supported by the government through the Ministry of Marine Affairs and Fisheries. In 2017, the santri actively participated in promoting the culture of fish consumption within the community. One of their approaches to enhance freshwater fish farming within the boarding school context was by fostering a culture of fish consumption. This initiative is expected to

contribute to increased protein consumption among the community, ultimately improving freshwater aquaculture based on the boarding school system [1]. The increasing consumer demand for catfish is being propelled by a growing awareness of the significance of fish consumption. This has led to the adoption of intensive farming practices to meet consumer demand for catfish. Intensive farming technology involves high stocking densities, aiming to produce yields that can meet the market demand for catfish. Feed plays a crucial role in intensive catfish farming. Commercial feed pellets significantly contribute to increasing production, but the high cost of these pellets poses a major obstacle in catfish farming. Approximately 60-70% of the production costs are allocated to feed expenses. The impact of intensive catfish farming is the accumulation of leftover feed and metabolic waste in the cultivation water, leading to a decrease in the water quality. This deterioration can even result in catfish mortality due to the build up of ammonia, nitrate, and nitrite. To ensure that feed can have a maximum impact, generate a larger biomass, reduce feed costs, and enhance the water quality in catfish farming, a solution needs to be found by incorporating probiotics in both feed and the cultivation water [2],[3].

Probiotics are living microorganisms suspended in a carrier medium that provide significant benefits to livestock by creating an ideal digestive environment and enhancing feed conversion efficiency. They make it easier for catfish to absorb the nutritional elements in their feed, promote livestock health, expedite growth, and offer protection against diseases. Probiotics can be administered through various means, including incorporation into feed, addition to water, or introduction through living intermediaries such as rotifers or artemia [4]. These probiotics produce enzymes like amylase, lipase, protease, and cellulase, which play a crucial role in breaking down nutrients like carbohydrates, proteins, and fats into simpler molecules. Fermentation of feed has been demonstrated to reduce the FCR value to as low as 0.997 in the cultivation of dumbo catfish fed with commercially fermented feed using probiofish [5]. The utilization of probiotics in feed at a dosage of 6 ml/kg resulted in the highest daily growth rate and feed efficiency, which were recorded at 3.12% and 31.65%, respectively [6].

Intensive cultivation of catfish (*Clarias gariepinus*) can lead to a decline in the quality of the cultivation medium, including a decrease in dissolved oxygen content and an increase in waste content, especially organic nitrogen [7]. Furthermore, the application of probiotics in feed, plays a significant role in improving water quality, enhancing productivity and feed efficiency, and reducing feed usage. The objective of this study is to investigate whether the addition of probiotics containing herbal ingredients to the fish feed can enhance the growth rate (SGR) and feed efficiency (FCR) of catfish.

26

2 Materials and methods

This research was conducted from June to July 2023 at the fish pond of PT Tirta Lestari Indonesia. The methodology employed in this study was an experimental approach using a Completely Randomized Design (CRD) with one factor having four treatment levels and three replications. The treatments applied in this research were as follows:

P0: Without probiotic supplementation

P1: Addition of probiotic at 2 cc/kg of feed

P2: Addition of probiotic at 4 cc/kg of feed

P3: Addition of probiotic at 6 cc/kg of feed

22

The catfish are fed three times a day, namely in the morning, noon, and evening until they are satiated. Length and weight measurements (sampling) are conducted every seven days. The research subject comprises catfish fingerlings ranging in size from 5 to 6, all derived from a single parent source purchased from a Hatchery located in Jombang, Jawa Timur, known for its reliable fingerling quality. The primary parameters measured include absolute weight, absolute length, specific growth rate, survival rate, feed efficiency, and feed conversion. Meanwhile, supporting parameters include water quality, such as temperature and pH. The sampling technique was performed by selecting 3 catfish from each pond, followed by weighing to determine the catfish's growth rate. Weighing was carried out under wet conditions by placing the catfish in another container filled with water of known weight. The difference in weight between the container with catfish and the container without catfish represented the individual weight of the catfish.

Data collection was conducted using an experimental method to examine the daily growth rate of catfish due to the administration of probiotic levels at 2 cc/kg of feed, 4 cc/kg of feed, 6 cc/kg of feed. This involved weighing the initial body weight of the research subjects and determining the total number of fingerlings. Subsequently, regular weighing was performed every week for a duration of 2 months. At the end of the study, the body weight was measured en masse, and the total number of fish was counted. The initial and final weighing were used to calculate the specific growth rate (SGR). Evaluating feed efficiency involved measuring the total feed provided to the catfish during the study and weighing the total initial and final biomass (FCR). Water quality in the catfish cultivation system was assessed using digital instruments, including a water temperature and pH meter. A TDS (Total Dissolved Solids) meter is utilized to quantify the total amount of dissolved substances in water. This encompasses various inorganic compounds and other dissolved substances found in water, such as minerals, salts, metals, ions, and more. TDS measurement can offer crucial insights into the level of contamination or the presence of compounds that can impact water quality and environmental health.

32

The collected data, which consists of primary parameters, is tabulated and analyzed using the SPSS application. This analysis involves employing Analysis of Variance (ANOVA) within a 95% confidence interval to determine whether the treatments significantly affect growth rate, feed conversion, and survival rate.

3 Results and discussion

2

3.1 Absolute weight, absolute length, specific growth rate, survival rate, feed efficiency, and feed conversion

12

Based on the observations of the average weight and length growth of catfish, there were noticeable differences in both length and weight between the probiotic-fermented feed treatment and the non-fermented feed treatment. The use of probiotic-fermented feed resulted in higher average length and weight compared to the control feed. The measurements for Specific Growth Rate (SGR), absolute weight, absolute length, survival rate, feed efficiency, and feed conversion of catfish are presented in Table 1.

Table 1. Specific Growth Rate (SGR), absolute weight, absolute length, survival rate, feed efficiency, and catfish feed conversion ratio.

Parameter	Treatment			
	P0	P1	P2	P3
absolute weight (g)	81,78±0,02 ^a	88,13±0,30 ^a	96,48±0,90 ^b	112,86±0,81 ^c
Absolute length (cm)	15,1±0,37 ^a	15,92±0,29 ^b	16,56±0,26 ^c	16,94±0,13 ^d
Specific Growth Rate (SGR) (%)	2,2271±0,046 ^a	2,3970±0,0605 ^b	2,419±0,009 ^c	2,4475±0,012 ^a
survival rate (%)	95,00±6,06	96,54±1,23	96,75±2,38	97,58±1,56
feed efficiency (%)	104,56±2,13 ^a	108,16±2,23 ^b	110,90±1,03 ^b	117,33±1,37 ^c
Feed Conversion Ratio (FCR)	¹⁴ 0,96±0,02 ^c	0,92±0,01 ^b	0,91±0,01 ^b	0,85±0,01 ^c

The highest absolute weight of catfish was observed in P3, which is the feed fermentation with a probiotic dosage of 6 ml/kg, measuring 112.86 grams, while the lowest was in the control treatment, which recorded 81.78 grams. The significance value in the ANOVA test is 0.000, indicating that $P < 0.05$, meaning that feed fermentation with probiotics has a significant effect on the absolute weight of catfish.

Fermented feed is more easily digested by fish compared to non-fermented feed, resulting in fish requiring less energy for digestion. The surplus energy can be utilized for various purposes, including fish growth, such as an increase in fish weight. After fermentation, most of the components are converted into simple compounds that can be directly used as fish feed, eliminating the need for further digestion by the fish [8]. The highest absolute length of catfish was obtained in treatment P3, measuring 16.94 cm, while the lowest was in P0, which recorded 15.1 cm. The results of the ANOVA test indicate that $P < 0.05$, signifying that feed fermentation with probiotics influences the absolute length of catfish. The growth in fish length is directly related to the growth in fish weight, which is why the highest absolute length was observed in P3, as the highest absolute weight was also achieved in P3.

The specific growth rate (SGR) of catfish, monitored every 7 days for 8 weeks following treatment, reveals that the highest SGR value was obtained through the application of a 6 cc/kg probiotic concentration, yielding an average SGR of 2,4475±0,012. In contrast, the lowest SGR was observed in the absence of probiotics (control), registering at 2,2271±0,046. The administration of probiotics containing Bio T₂₀, which are heterotrophic bacteria, in different quantities through both food and water has a significant impact on the specific growth rate (SGR) of catfish fry raised for 8 weeks, with statistically significant differences observed ($P < 0.05$). This suggests that catfish are capable of utilizing fermented feed for growth during their cultivation. The catfish's growth is clearly reflected in the increases in body weight and specific growth rate (SGR) values over the 8-week cultivation period. The

specific growth rate of catfish exceeds that of previous research involving the supplementation of probiotics to catfish. For instance, introducing probiotics at a concentration of 108 CFU/mL into the Dumbo catfish feed yielded a specific growth rate of 2.16% [9].

The survival rate in this study ranges from 95.00% to 97.58%. It is believed that this phenomenon is a result of probiotic bacteria activity, particularly *Lactobacillus* sp. These bacteria have the capability to generate lactic acid from sugars and other carbohydrates produced by photosynthetic bacteria and yeast. *Lactobacillus* sp. bacteria contribute to the equilibrium of the gut microbiota, thereby enhancing fish digestion through the conversion of carbohydrates into lactic acid, leading to a reduction in pH. Consequently, this process stimulates the production of naturally occurring enzymes, which improves nutrient absorption, feed intake, growth, and acts as a deterrent to pathogenic organisms [10]. The utilization of probiotic technology has led to a notably high survival rate among catfish, as it effectively preserves the quality of the aquatic environment.

The efficiency of catfish feeding is remarkably high. The peak feed efficiency value was achieved in P3, reaching 117.33%, whereas the lowest was documented in P0 at 104.56%. This is the reason behind the notably high feed efficiency observed in this research. The addition of carbohydrates to the cultivation medium can be converted into an energy source by heterotrophic bacteria leading to the production of a significant quantity of protein-rich bacterial biomass. This bacterial biomass can be subsequently utilized by fish as a plentiful source of high-protein supplementary nutrition [11].

The lower feed conversion ratio in P3 is believed to be due not only to the fact that fish utilize the flocculants for food but also because the fermentation process results in higher fish absorption of the feed. The digestive enzymes produced by microbes during the fermentation process assist in breaking down complex compounds into simpler components, making the feed more easily absorbed by the intestines [12].

The feed conversion values for all treatments are considered favorable, ranging from 0.96 to 0.85. Feed Conversion Ratio (FCR) is the ratio of the amount of feed required to produce 1 kg of fish meat. The FCR of catfish in the probiotic treatment is lower compared to the control group. Feed utilization efficiency signifies the amount of feed that can be utilized by the fish, converting it into an increase in the fish's body weight. Feed efficiency can be assessed through various factors, with one of them being the FCR. The highest level of feed utilization efficiency is achieved when the calculation of the feed conversion ratio is the lowest, indicating superior feed quality compared to other treatments [13]. Good feed quality results in the catfish channeling more energy towards growth, thus anticipating improved growth rates with reduced feed intake. Factors influencing the level of feed efficiency include the type and quantity of each nutrient source in the feed. The quantity and quality of the feed provided to the catfish significantly affect their growth. The higher the feed efficiency value, the more favorable the catfish's response to the feed, as evidenced by rapid growth rates.

4 Conclusion

The research findings indicate that the provision of fermented feed has an impact on body weight growth, absolute length, feed efficiency, and feed conversion, but it does not affect the survival rate of catfish. The best results were obtained in P3, which involved feed fermentation with a probiotic dosage of 6 cc/kg of feed, resulting in a specific growth rate of 2.44%, an absolute weight of 112.86 grams, an absolute length of 15.51 cm, a survival rate of 97.58%, feed efficiency of 117.33%, and a feed conversion of 0.85.

Acknowledgements

We acknowledge the support received from Universitas Pesantren Tinggi Darul Ulum for the support, PT Tirta Lestari Indonesia, Directorate General of Research and Higher Education, Ministry of Education and Culture, Republic of Indonesia, for their generous funding support through the Matching Fund Program in 2023.

Reference

- [1] Republika, Gus Sholah Dorong Pesantren Membudidayakan Lele. 2017. <https://khazanah.republika.co.id/berita/otutph384/gus-sholah-dorong-pesantren-membudidayakan-lele>
- [2] Arief, M., Fitriani, N. dan Subekti, S. . 2014. Pengaruh pemberian probiotik berbeda pada pakan komersial terhadap pertumbuhan dan efisiensi pakan lele Sangkuriang (*Clarias sp.*). Jurnal Ilmiah Perikanan dan Kelautan, . 6(1): 49-53. April 2014.
- [3] Gunawan, R.G.B dan Bagus. H. 2011. Dongkrak produksi lele dengan probiotik organik. Agromedia Pustaka. Jakarta.
- [4] Irianto, A. Potensi mikroorganisma : di atas langit ada langit. Ringkasan Orasi ilmiah di fakultas Biologi Universitas Jenderal Sudirman tanggal 12 mei, 2007.
- [5] Negara I. K. W., Marsoedi dan Edi S. 2015. Strategi pengembangan budidaya lele dumbo *Clarias sp.* melalui Program pengembangan usaha mina pedesaan perikanan budidaya Di kabupaten buleleng. J. Manusia dan lingkungan. 22 (3): 365-371
- [6] Ahmadi, H., Iskandar, Kurniawati, N., Pemberian Probiotik Dalam Pakan Terhadap Pertumbuhan LEle Sangkuriang (*Clarias gariepinus*) Pada Pendederan II, Jurnal Perikanan dan Kelautan, Vol. 3., No. 4., Hal 99-107. 2012
- [7] Rachmawati, D., I. Samidjan. dan H.Setyono. 2015. Manajemen kualitas air media budiaya ikan lele Sangkuriang (*Clarias gariepinus*) dengan teknik probiotik pada kolam terpal di desa Voksi Rektosari, Kecamatan Suruh, Kabupaen Semarang. PENA Akuatika. Volume 12 No. 1 - September 2015.
- [8] Purnomo, P. D. 2012. Pengaruh Penambahan Karbohidrat Pada Media Pemeliharaan Terhadap Produksi Budidaya Intensif Nila (*Oreochromis niloticus*). Journal of Aquaculture Management and Technology. 1 (1): 161-179.
- [9] Wardika A. S., Suminto, dan Agung S. 2014. Pengaruh Bakteri Probiotik Pada Pakan dengan Dosis Berbeda Terhadap Efisiensi Pemanfaatan Pakan, Pertumbuhan dan Kelulushidupan Lele Dumbo p(*Clarias gariepinus*). Journal of Aquaculture Management and Technology, 3 (4) : 9-17 hal
- [10] Arief, M., Pemberian Probiotik yang Berbeda pada Pakan Komersil terhadap Pertumbuhan Retensi Protein dan Serat Kasar pada Ikan Nila (*Oreochromis sp.*). Argoveteriner. 1 (2): 88 – 93. 2013.
- [11] Purnomo, P. D., Pengaruh Penambahan Karbohidrat Pada Media Pemeliharaan Terhadap Produksi Budidaya Intensif Nila (*Oreochromis niloticus*). Journal of Aquaculture Management and Technology. 1 (1): 161-179. 2012

- [12] Sugih F.H. 2005. Pengaruh penambahan probiotik dalam pakan komersil terhadap pertumbuhan benih ikan gurami (*Osphronemus goramy Lac.*). [Skripsi]. Jurusan Perikanan, Unpad. Bandung.
- [13] Hastuti dan Subandiyono. 2014. Performa Produksi Ikan Lele Dumbo (*Clarias gariepinus, Burch*) Yang Dipelihara Dengan Teknologi Biofloc. *Jurnal Saintek Perikanan*, 10 (1) : 37-42 hal.

ORIGINALITY REPORT

20%

SIMILARITY INDEX

12%

INTERNET SOURCES

15%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	Deswati Deswati, Rahmiana Zein, Suparno Suparno, Hilfi Pardi. "Modified biofloc technology and its effects on water quality and growth of catfish", Separation Science and Technology, 2023 Publication	2%
2	garuda.ristekbrin.go.id Internet Source	2%
3	RC Mukti, M Amin, Yulisman. "Utilization of alternative ingredients as combination feed on the growth of catfish (Clarias sp.)", IOP Conference Series: Earth and Environmental Science, 2021 Publication	2%
4	www.tandfonline.com Internet Source	1%
5	www.globalscientificjournal.com Internet Source	1%
6	S N Pratama, R C Mukti. "Utilization of herbal supplements on feed on growth and survival rate of catfish (Clarias sp.)", IOP Conference Series: Earth and Environmental Science, 2021 Publication	1%
7	ijariie.com Internet Source	1%
8	www.neliti.com Internet Source	1%

9	E Aristasari, R A Nur 'Aini, W Nopita, Agustono, M Lamid, M A Al-Arif. "The growth, protein content, and fatty acid of catfish meat (pangasius sp.) With the addition of different lysine doses in commercial feed", IOP Conference Series: Earth and Environmental Science, 2020 Publication	1 %
10	www.mdpi.com Internet Source	1 %
11	www.jatit.org Internet Source	1 %
12	Widowati Budijastuti, Nur Ducha, Dyah Hariani, Raharjo. "The Effect of Giving Worm Manure on The Growth of Catfish and Kale in Buckets", E3S Web of Conferences, 2021 Publication	<1 %
13	Submitted to Adtalem Global Education, Inc. Student Paper	<1 %
14	K. M. HAVSTAD. "GROWTH PATTERNS OF RANGE BEEF CALVES OVER DISCRETE PREWEANING INTERVALS", Canadian Journal of Animal Science, 12/1989 Publication	<1 %
15	www.bioflux.com.ro Internet Source	<1 %
16	docobook.com Internet Source	<1 %
17	I Putra, I Efendi, I Lukisyowati, U M Tang, R Rusliadi, R A Manik. " Effects of Various Carbon Doses on Tilapia (sp) Culture with Biofloc Technology ", IOP Conference Series: Earth and Environmental Science, 2020 Publication	<1 %

18	terubuk.ejournal.unri.ac.id Internet Source	<1 %
19	www.jurnal.unsyiah.ac.id Internet Source	<1 %
20	Ita Apriani, Elsari Tanjung Putri. "PENGARUH PROBIOTIK PADA PAKAN TERHADAP PERTUMBUHAN DAN KELANGSUNGAN HIDUP IKAN LELE MUTIARA (<i>Clarias gariepinus</i>) BUDIDAYA SISTEM BIOFLOK", Jurnal Ruaya : Jurnal Penelitian dan Kajian Ilmu Perikanan dan Kelautan, 2021 Publication	<1 %
21	M M El Feky. "Evaluation the Effect of Local and Imported Yeasts as Supplementary Food on the African Catfish (<i>Clarias gariepinus</i>) in Egypt", Journal of Aquaculture & Marine Biology, 2015 Publication	<1 %
22	Robert Devi Anugrah, Rafvenia, Meitiyani, Luthpi Safahi. "The Effect of Eggshell Organic Fertilizer on Vegetative Growth of Cayenne Pepper (<i>Capsicum frutescens</i> L)", IOP Conference Series: Earth and Environmental Science, 2021 Publication	<1 %
23	core.ac.uk Internet Source	<1 %
24	ejournal3.undip.ac.id Internet Source	<1 %
25	www.scilit.net Internet Source	<1 %
26	"Proceedings of the Andalas International Public Health Conference 2017", BMC Public Health, 2017 Publication	<1 %

27 123dok.com <1 %
Internet Source

28 Abdul Rakhfid, Wa Ode Nanny Kulsum, Fendi Fendi, Mosriula Mosriula et al. "The use of probiotic for growth and survival of milkfish (*Chanos chanos* Forskal)", *Akuatikisile: Jurnal Akuakultur, Pesisir dan Pulau-Pulau Kecil*, 2020 <1 %
Publication

29 Mehmet Hayta, Elif Meltem İşçimen. "Effects of Phytochemical Fortification of Flour and Bread on Human Health", Elsevier BV, 2019 <1 %
Publication

30 jurnal.faperta.untad.ac.id <1 %
Internet Source

31 seminarfkp.undana.ac.id <1 %
Internet Source

32 www.atlantis-press.com <1 %
Internet Source

Exclude quotes Off

Exclude matches Off

Exclude bibliography On